



ORAL SURGERY ASSOCIATES

**Dr. R. Brent Bailey, D.D.S. Dr. Michael J. Anton, D.D.S. Dr. Garrett J. Seeba, D.M.D.,
M.D.**

595 E. Medical Center Boulevard
Webster, TX 77598

(281) 461-1982

AFTER TOOTH EXTRACTION

After tooth extraction, it is important for a blood clot to form to stop the bleeding and begin the healing process. That is why we ask you to bite on a gauze pad for 45-60 minutes after the extraction. If the bleeding or oozing still persists, place another gauze pad and bite firmly for another 30 minutes. You may have to do this several times.

After the blood clot forms, it is important not to disturb or dislodge the clot, as it aids healing. Do not rinse vigorously, suck on straws, smoke, drink alcohol or brush teeth next to the extraction site for 72 hours. These activities will dislodge or dissolve the clot and retard the healing process. Limit vigorous exercise for the next 24 hours as exercise will increase blood pressure and may cause more bleeding from the extraction site.

After the tooth is extracted, you may feel some pain and may experience some swelling. An ice pack or an unopened bag of frozen peas or corn applied to the area will keep swelling to a minimum. Place a cloth between the ice pack and your face. Take pain medications as prescribed. The swelling usually subsides after 48 hours.

Use the pain medication as directed. Call the office if the medication does not seem to be working. If antibiotics are prescribed, continue to take them for the indicated length of time, even if signs and symptoms of infection are gone. Drink lots of fluid and eat nutritious soft food on the day of the extraction. You can eat normally as soon as you are comfortable.

It is important to resume your normal dental routine after 24 hours. This routine should include brushing and flossing your teeth at least once a day. This regimen will speed healing.

After a few days you will feel fine and can resume your normal activities. If you have heavy bleeding, severe pain, continued swelling for 2-3 days, or a reaction to the medication, call our office immediately at 281-461-1982.